

A fast algorithm to estimate the cost and the right-hand side parameter vectors in inverse linear optimization

EURO 2019, Dublin, Ireland

Juan M. Morales¹, Ricardo Fernández-Blanco¹, Salvador Pineda¹

¹OASYS Research Group, University of Malaga, Spain

June 25, 2019



European Research Council

Established by the European Commission



Subject of Study: Inverse linear optimization

$$\begin{array}{ll} \underset{\mathbf{x}}{\text{Minimize}} & (\mathbf{c}^T + \mathbf{s}^T)\mathbf{x} \\ \text{Subject to} & \mathbf{A}\mathbf{x} \leq \mathbf{b} \end{array}$$


Objective: Estimate $\mathbf{c}$ and $\mathbf{b}$ from data (observations) $\left\{(\tilde{\mathbf{x}}_t, \mathbf{s}_t)_{t=1}^T\right\}$, i.e.,

Input: $\{\mathbf{s}_t, t = 1, \dots, T\}$

Output: $\{\mathbf{x}_t, t = 1, \dots, T\}$, but we observe $\{\tilde{\mathbf{x}}_t, t = 1, \dots, T\}$, as measurements may be contaminated with *noise*

Motivating example¹

$$\begin{aligned}
 \max_{x_b, \forall b} \quad & \sum_{b=1}^B x_b (u_b - p) \\
 \text{s. t.} \quad & \underline{P} \leq \sum_{b=1}^B x_b \leq \overline{P} \quad (\underline{\lambda}, \overline{\lambda}) \\
 & 0 \leq x_b \leq E_b \quad (\underline{\phi}_b, \overline{\phi}_b)
 \end{aligned}$$



Objective: Estimate u_b , $b = 1, \dots, B$, $\underline{P}$ and $\overline{P}$ from observations $\{(\tilde{x}_t, p_t)_{t=1}^T\}$, i.e.,

Input: $\{p_t, t = 1, \dots, T\}$

Output: $\{x_{b,t}, b = 1, \dots, B; t = 1, \dots, T\}$, but we only observe the (noisy) *aggregate* consumption $\{\tilde{x}_t, t = 1, \dots, T\}$

¹J. Saez-Gallego, J. M. Morales, M. Zugno and H. Madsen (2016). A data-driven bidding model for a cluster of price-responsive consumers of electricity. *IEEE Transactions on Power Systems* 31(6): 5001–5011

Motivating example

Time-variant
parameters

$$\underline{P}_t = \underline{\mu} + \sum_{r=1}^R \underline{\alpha}_r Z_{r,t}$$

$$\overline{P}_t = \overline{\mu} + \sum_{r=1}^R \overline{\alpha}_r Z_{r,t}$$

$$u_{b,t} = \mu_b^u + \sum_{r=1}^R \alpha_r^u Z_{r,t}$$

Objective: Estimate $\underline{\mu}$, $\overline{\mu}$; μ_b^u , $b = 1, \dots, B$; $\underline{\alpha}_r$, $\overline{\alpha}_r$, and α_r^u , $r = 1, \dots, R$ from observations $\left\{ (\tilde{x}_t, p_t, Z_{1,t}, Z_{2,t}, \dots, Z_{R,t})_{t=1}^T \right\}$

Estimation: Duality-gap minimization

$$\begin{aligned}
 & \underset{\Omega}{\text{minimize}} && \sum_{t=1}^T \epsilon_t \\
 & \text{subject to} && \bar{P}_t \bar{\lambda}_t - \underline{P}_t \underline{\lambda}_t + \sum_{b=1}^B E_b \bar{\phi}_{b,t} - \epsilon_t = \sum_{b=1}^B x_{b,t} (u_{b,t} - p_t), \quad \forall t \\
 & && \bar{\phi}_{b,t} - \underline{\phi}_{b,t} + \bar{\lambda}_t - \underline{\lambda}_t = u_{b,t} - p_t, \quad \forall t \\
 & && \bar{\phi}_{b,t}, \underline{\phi}_{b,t}, \bar{\lambda}_t, \underline{\lambda}_t \geq 0, \quad \forall t \\
 & && \epsilon_t \geq 0, \quad \forall t
 \end{aligned}$$

$$\Omega = \left\{ \left(\epsilon_t, \bar{P}_t, \underline{P}_t, u_{b,t}, \bar{\lambda}_t, \underline{\lambda}_t, \bar{\phi}_{b,t}, \underline{\phi}_{b,t} \right)_{t=1}^T \right\}$$

Estimation: Duality-gap minimization²

- The estimation problem is **non-linear and non-convex**.
- We statistically **approximate its solution** by solving two **linear programming problems** instead.
 - ① A **feasibility** problem (estimation of power bounds).
 - ② An **optimality** problem (estimation of marginal utilities).
- A **two-step data-driven estimation procedure** to achieve **optimality** and **feasibility** of $\tilde{\mathbf{x}}$ in a statistical sense.

²J. Saez-Gallego and J. M. Morales (2018). Short-term forecasting of price-responsive loads using inverse optimization. *IEEE Transactions on Smart Grid* 9(5):4805–4814

Feasibility problem: Estimation of power bounds

$$\text{Minimize}_{\underline{P}, \bar{P}, \xi, \mu, \alpha} \sum_{t=1}^T \left((1-K) (\bar{\xi}_t^+ + \underline{\xi}_t^+) + K (\bar{\xi}_t^- + \underline{\xi}_t^-) \right)$$

subject to

$$\bar{P}_t - x'_t = \bar{\xi}_t^+ - \bar{\xi}_t^- \quad \forall t$$


$$x'_t - \underline{P}_t = \underline{\xi}_t^+ - \underline{\xi}_t^- \quad \forall t$$

$$\underline{P}_t \leq \bar{P}_t \quad \forall t$$

$$\underline{P}_t = \underline{\mu} + \sum_{r=1}^R \underline{\alpha}_r Z_{r,t} \quad \forall t$$

$$\bar{P}_t = \bar{\mu} + \sum_{r=1}^R \bar{\alpha}_r Z_{r,t} \quad \forall t$$

$$0 \leq \bar{\xi}_t^+, \bar{\xi}_t^-, \underline{\xi}_t^+, \underline{\xi}_t^- \quad \forall t$$



$$\hat{\bar{P}}_t, \hat{\underline{P}}_t, \hat{\underline{\mu}}, \hat{\bar{\mu}}, \hat{\underline{\alpha}}_r, \hat{\bar{\alpha}}_r$$

K controls the percentage of observations falling within $[\underline{P}, \bar{P}]$

Optimality problem: Estimating marginal utilities

$$\text{Minimize}_{\Omega} \sum_{t=1}^T \epsilon_t$$

$$\text{subject to } \hat{\bar{P}}_t \bar{\lambda}_t - \hat{\underline{P}}_t \underline{\lambda}_t + \sum_{b=1}^B E_b \bar{\phi}_{b,t} - \epsilon_t =$$

$$\sum_{b=1}^B \tilde{x}'_{b,t} (u_{b,t} - p_t)$$

$$\forall t$$

$$-\underline{\phi}_{b,t} + \bar{\phi}_{b,t} - \underline{\lambda}_t + \bar{\lambda}_t = u_{b,t} - p_t \quad \forall b, t$$

$$u_{b,t} = \mu_b^u + \sum_r \alpha_r^u Z_{r,t} \quad \forall b, t$$

$$\mu_b^u \geq \mu_{b+1}^u \quad \forall b < B$$

$$\mu_1^u \geq 200 + \mu_2^u$$

$$0 \leq \bar{\lambda}_t, \underline{\lambda}_t, \underline{\phi}_{b,t}, \bar{\phi}_{b,t}, \epsilon_t \quad \forall b, t.$$




x' optimal?

$$\hat{u}_{b,t}, \hat{\mu}_b^u, \hat{\alpha}_r^u$$

Solving the estimation problem

In the **bound estimation problem**, the **penalty parameter K** is statistically tuned through **validation**:



Noiseless case: $\tilde{X}_t = X_t, \quad t = 1, \dots, T$


- We assume $p_t \sim N(0.050, 0.0125)$; $\mathbf{Z} = (p_{t-1}, p_{t-2})$
- We analyze 3 cases:
 - Case 1: $(u_{b,t}, \underline{P}_t, \overline{P}_t)$ are time-invariant
 - Case 2: $u_{b,t}$ is time-variant and $(\underline{P}_t, \overline{P}_t)$ are time-invariant
 - Case 3: $(u_{b,t}, \underline{P}_t, \overline{P}_t)$ are time-variant
- Let us assume $u_{1,t} = 0.2, \mu_2^u = 0.070, \mu_b^u = \mu_{b-1} - 0.0015 \quad \forall b > 2, \underline{\mu} = 0, \overline{\mu} = 100$

Case	α_r^u		$\underline{\alpha}_r$		$\overline{\alpha}_r$	
	1	2	1	2	1	2
1	0.0	0.0	0.0	0.0	0.0	0.0
2	-0.5	0.3	0.0	0.0	0.0	0.0
3	-0.5	0.3	100	-20	200	-100

- We consider 8 data samples of sizes: 150, 200, 300, 400, 500, 1000, 2000, 3000. $\Omega^{test} = 100$ last periods (identical for all of them); $\Omega^{tr} / \Omega^v \Rightarrow 70\%/30\%$

Noiseless case: $\tilde{X}_t = X_t, \quad t = 1, \dots, T$

- Normalized root mean square error





- Optimal hyper-parameter K tends to 1 when increasing the sample size
- Estimated parameter for case 1 (time-invariant!)

Sample size	μ_2^U	$\underline{\mu}$	$\overline{\mu}$
150	0.062	21.05	100.0
200	0.067	5.26	100.0
300	0.070	0.00	100.0
True	0.070	0.00	100.0



Noiseless case: $\tilde{X}_t = X_t, \quad t = 1, \dots, T$

- Results for case 2: $u_{b,t}$ is time-variant and $(P_t, \bar{P}_t)$ are time-invariant



Noiseless case: $\tilde{x}_t = x_t, \quad t = 1, \dots, T$

- Results for case 3: $(u_{b,t}, \underline{P}_t, \overline{P}_t)$ are time-variant



Noiseless case: $\tilde{x}_t = x_t, \quad t = 1, \dots, T$

Takeaway message: Our two-step estimation procedure performs satisfactorily in a noiseless setup, but...

What if the observations $(\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_T)$ are corrupted with some measurement i.i.d. noise $\omega_t \sim N(0, \sigma)$?

Noisy case: $\tilde{x}_t = x_t + \omega_t, \quad t = 1, \dots, T; \quad \omega_t \sim N(0, \sigma)$

Consider the following illustrative linear program³:

$$\begin{aligned} \min_x \quad & (\theta + p) \cdot x \\ \text{s. t.} \quad & -1 \leq x \leq 1 \end{aligned}$$

Objective: Estimate θ from observations $(\tilde{x}_t, p_t)$ where:

$p_1, p_2, \dots, p_t, \dots, p_T$ i.i.d, with $p_t \sim U(-2, 1)$


$\tilde{x}_t = x_t + \omega_t$, with $\omega_1, \dots, \omega_t, \dots, \omega_T$ i.i.d with $\omega_t \sim N(0, 1)$

³A. Aswani, Z.-J. Shen and A. Siddiq (2018). Inverse optimization with noisy data. *Operations Research* 66(3):870–892

Noisy case: $\tilde{x}_t = x_t + \omega_t$, $t = 1, \dots, T$; $\omega_t \sim N(0, \sigma)$

Optimality problem:

$$\begin{aligned}
 & \min_{\hat{\theta}, \epsilon_t, \bar{\beta}_t, \underline{\beta}_t} \sum_{t=1}^T \epsilon_t \\
 & \text{s. t. } (\hat{\theta} + p_t) \cdot \tilde{x}_t - \epsilon_t = -\bar{\beta}_t - \underline{\beta}_t, \quad \forall t \\
 & \quad \underline{\beta}_t - \bar{\beta}_t = \theta + u_t, \quad \forall t \\
 & \quad \bar{\beta}_t, \underline{\beta}_t \geq 0, \quad \forall t \\
 & \quad \epsilon_t \geq 0, \quad \forall t
 \end{aligned}$$



Unable to recover the true parameters!

Noisy case: $\tilde{x}_t = x_t + \omega_t$, $t = 1, \dots, T$; $\omega_t \sim N(0, \sigma)$

Any idea to preserve statistical consistency?

Let us assume **asymptotic regime**, i.e., $T \uparrow\uparrow$:

Weak duality: $(\hat{\theta} + p_t) \cdot x_t \geq -\bar{\beta}_t - \underline{\beta}_t$

Strong duality: $(\theta + p_t) \cdot x_t = -\bar{\beta}_t(\theta) - \underline{\beta}_t(\theta)$

$$(\hat{\theta} + p_t) \cdot \tilde{x}_t - \epsilon_t = -\bar{\beta}_t - \underline{\beta}_t \Rightarrow (\hat{\theta} + p_t) \cdot (x_t + \omega_t) - \epsilon_t = -\bar{\beta}_t - \underline{\beta}_t \Rightarrow$$

$$\Rightarrow (\hat{\theta} + p_t) \cdot x_t + \hat{\theta}\omega_t + p_t\omega_t - \epsilon_t = -\bar{\beta}_t - \underline{\beta}_t$$


Since $\frac{1}{T} \sum_{t=1}^T \omega_t \rightarrow 0$ and $\frac{1}{T} \sum_{t=1}^T p_t \omega_t \rightarrow 0$ when $T \uparrow\uparrow$, we have

$$\sum_{t=1}^T \epsilon_t = \sum_{t=1}^T \left((\hat{\theta} + p_t) \cdot x_t + \bar{\beta}_t + \underline{\beta}_t \right) \geq 0 \text{ and } = 0 \text{ iff } \hat{\theta} = \theta$$

Noisy case: $\tilde{x}_t = x_t + \omega_t$, $t = 1, \dots, T$; $\omega_t \sim N(0, \sigma)$


Recast optimality problem

$$\begin{aligned}
 & \min_{\theta, \epsilon_t, \bar{\beta}_t, \underline{\beta}_t} \sum_{t=1}^T \epsilon_t \\
 \text{s. t. } & (\theta + p_t) \cdot \tilde{x}_t - \epsilon_t = -\bar{\beta}_t - \underline{\beta}_t, \quad \forall t \\
 & \underline{\beta}_t - \bar{\beta}_t = \theta + u_t, \quad \forall t \\
 & \bar{\beta}_t, \underline{\beta}_t \geq 0, \quad \forall t \\
 & \epsilon_t \geq 0, \quad \forall t : \tilde{x}_t \in [-1, 1]
 \end{aligned}$$



Minimizing $\left| \sum_{t=1}^T \epsilon_i \right|$ or $\left(\sum_{t=1}^T \epsilon_i \right)^2$ should be theoretically equivalent when $T \uparrow$, but not in practice (finite-sample regime)

Noisy case: $\tilde{x}_t = x_t + \omega_t$, $t = 1, \dots, T$; $\omega_t \sim N(0, \sigma)$




Noisy case: $\tilde{x}_t = x_t + \omega_t, \quad t = 1, \dots, T; \quad \omega_t \sim N(0, \sigma)$

Statistical consistency is a theoretical nice property, but not that useful in practice, where we normally work under a **finite-sample regime** (that is, T not big enough)


What about the **bias** and **variance** of the estimated parameters?

Noisy case: $\tilde{x}_t = x_t + \omega_t$, $t = 1, \dots, T$; $\omega_t \sim N(0, \sigma)$



Checking for **bias**: Estimator average over 100 instances of $T = 500$


Noisy case: $\tilde{x}_t = x_t + \omega_t$, $t = 1, \dots, T$; $\omega_t \sim N(0, \sigma)$



Checking for **bias** and **variance**: Box plot of the parameter estimator computed for 500 instances of $T = 200$ (true value $\theta = 0$)

Noisy case: $\tilde{x}_t = x_t + \omega_t, \quad t = 1, \dots, T; \quad \omega_t \sim N(0, \sigma)$

Bias-variance trade-off: Minimize $L \sum_t^T |\epsilon_t| + (1 - L) \left| \sum_{t=1}^T \epsilon_i \right|$, with L being a control parameter ($L \rightarrow 0$ as $T \uparrow$)



Conclusions

- A fast method for inverse linear optimization able to recover the true parameters under a noiseless setup (even in the case of unknown right-hand side parameter vector)
- Different alternatives to preserve statistical consistency in the presence of noise in the measurements
- These alternatives exhibit different behaviors in terms of bias and variance

Future Work

- Need for theoretical insight
- Extension to the case of unknown parameters in the constraints
- Extension to inverse convex optimization

Contacts

Any questions?



European Research Council

Established by the European Commission

Juan Miguel Morales

juan.morales@uma.es

OASYS Webpage: oasys.uma.es

